

Infusion

Blood Products Transfusion Order



PATIENT INFORMATION

Last Name: _____ First Name: _____ MI: _____ DOB: _____
Address: _____ Phone: _____
City: _____ State: _____ Zip Code: _____
Emergency Contact: _____ Relationship: _____ Phone: _____

PROVIDER INFORMATION

Referring Provider: _____ Phone Number: _____ Fax Number: _____

ADDITIONAL INFORMATION

Check if Patient is uninsured. Provide ICD-10 code and description: _____
Weight: _____ Allergies: _____
Is the patient ambulatory? Yes No Does the patient require bariatric equipment? Yes No

ORDERS PRECEDED BY A REQUIRE A TO INITIATE THE ORDER.)

Red blood Cell transfusion:

Type & Cross and hold **OR** Type & Cross and Transfuse _____ Units of PRBC (SH uses leukoreduced CMV safe RBCs)
Irradiate Unit? Yes NO (order will be returned if not selected)

Type & Cross and Transfuse:

(select only one time interval) STAT (Within 24 hrs) **OR** Within 25-48 hrs. **OR** Within 49-72 hrs.

Platelets:

Transfuse _____ Units Irradiate Unit? Yes No (order will be returned if not selected) HLA Cross match
(HLA testing is performed through American Red Cross and takes approx. 3 working days)
(select only one time interval) STAT (Within 24 hrs.) **OR** within 25-48 hrs. **OR** within 49-72 hrs.

Pre-Medications: Select One. NO Pre-Medications **OR** Administer pre-meds 30 min prior to transfusion

Acetaminophen P.O. Every 4 hrs. (select one dose) 500 mg **OR** 650 mg **OR** _____ mg
 Diphenhydramine (select one dose) 12.5mg IV **OR** 25mg IV **OR** 50mg IV
 Dexamethasone (select one dose) 4mg IV **OR** 8mg IV **OR** 10mg IV
 Furosemide IV (select dose and frequency) _____ mg IV (select one) _____ after each unit **OR** _____ only Once

Other Instructions:

HCT _____ Hgb _____ Plt _____ date of results _____

PATIENTS WITH CENTRAL LINE ACCESS

- Central line care per Salem Health CVAD Access Policy. (Lippincott) or routine implanted port care per manufacture device maintenance card if card is available.
- Alteplase per Salem Health Central Venous Access Device declotting (Lippincott) for S/sx of occlusion: Inability to infuse fluids, no blood return, increased resistance when flushing, increased occlusion/high-pressure alarm when using an infusion pump, sluggish gravity flow.
- View Chest X-ray to verify catheter tip location PRN for: Catheter migration greater than 5 cm (PICC only), signs and symptoms of tip malposition (occlusion unresolved by Cathflo, discomfort in the arm, neck or chest, unusual sensations or sounds when flushing, neck vein engorgement, or heart palpitations.) Notify Physician or Provider
- Blood bank may substitute irradiated blood or platelets for non-irradiated or Psorlan treated platelets for irradiated or non-irradiated based on availability per blood bank policy
- Follow SH OP Infusion reaction algorithm for symptoms of an infusion reaction. Notify provider if implemented. Follow steps defined in the SH blood administration for adults policy (section C 1-3)

Provider Signature

Provider Printed Name

Date: